

High Resolution Time Series Measurements of Bio-optical and Physical Variability in the Coastal Ocean as Part of HyCODE

T. Dickey
Ocean Physics Laboratory
University of California at Santa Barbara
6487 Calle Real, Suite A
Goleta, CA 93117
phone: (805) 893-7354 fax: (805) 967-5704 email: tommy.dickey@opl.ucsb.edu

Award #: N000149910222
<http://www.opl.ucsb.edu/hycode.html>

LONG-TERM GOALS

This project is intended to increase understanding of the variability of inherent and apparent optical properties (IOPs and AOPs) of ocean waters and their relationships to each other as well as to physical processes on continental shelves. Data collected during our study and their analyses also expand the utility of remotely sensed hyperspectral ocean color data.

OBJECTIVES

Specific objectives of the project are

1. To provide the maximum number of *in situ* observations (highest possible number of match-ups) of IOPs and AOPs possible for calibrating, groundtruthing, and relating subsurface optical properties (algorithm development) to aircraft and spacecraft ocean color data, and to develop, test, and validate optical models and high resolution interdisciplinary models of the coastal ocean.
2. To study processes which contribute to temporal and spatial (horizontal and vertical) variability of spectral IOPs and AOPs. We are determining how temporal and spatial variability in IOPs and AOPs are affected by:
 - a) Coastal physical and biological dynamics (upwelling/downwelling, fronts, filaments, eddies, blooms, etc.) and larger scale circulation patterns
 - b) Wave fields (e.g., tides and surface, internal, and solitary waves)
 - c) Water column stratification and current shears
 - d) Near surface and near bottom mixing (e.g., effects on primary productivity, sediment resuspension, dilution, dispersion, etc.)
 - e) Diurnal and seasonal biological and physical cycles

| Report Documentation Page | | | | Form Approved OMB No. 0704-0188 | |
|--|------------------------------------|-------------------------------------|---|---|---------------------------------|
| Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. | | | | | |
| 1. REPORT DATE 30 SEP 2002 | | 2. REPORT TYPE | | 3. DATES COVERED 00-00-2002 to 00-00-2002 | |
| 4. TITLE AND SUBTITLE High Resolution Time Series Measurements of Bio-optical and Physical Variability in the Coastal Ocean as Part of HyCODE | | | | 5a. CONTRACT NUMBER | |
| | | | | 5b. GRANT NUMBER | |
| | | | | 5c. PROGRAM ELEMENT NUMBER | |
| 6. AUTHOR(S) | | | | 5d. PROJECT NUMBER | |
| | | | | 5e. TASK NUMBER | |
| | | | | 5f. WORK UNIT NUMBER | |
| 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Ocean Physics Laboratory, University of California at Santa Barbara,,6487 Calle Real, Suite A,,Goleta,,CA, 93117 | | | | 8. PERFORMING ORGANIZATION REPORT NUMBER | |
| 9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) | | | | 10. SPONSOR/MONITOR'S ACRONYM(S) | |
| | | | | 11. SPONSOR/MONITOR'S REPORT NUMBER(S) | |
| 12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited | | | | | |
| 13. SUPPLEMENTARY NOTES | | | | | |
| 14. ABSTRACT This project is intended to increase understanding of the variability of inherent and apparent optical properties (IOPs and AOPs) of ocean waters and their relationships to each other as well as to physical processes on continental shelves. Data collected during our study and their analyses also expand the utility of remotely sensed hyperspectral ocean color data. | | | | | |
| 15. SUBJECT TERMS | | | | | |
| 16. SECURITY CLASSIFICATION OF: | | | 17. LIMITATION OF ABSTRACT Same as Report (SAR) | 18. NUMBER OF PAGES 7 | 19a. NAME OF RESPONSIBLE PERSON |
| a. REPORT unclassified | b. ABSTRACT unclassified | c. THIS PAGE unclassified | | | |

f) Riverine and runoff inflows (i.e., changes in relative contributions of sedimentary, biological, and colored dissolved material (CDM) components, buoyant plumes, and phytoplankton blooms).

APPROACH

Intensive HyCODE field observations were made in the summers of 2000 and 2001. The particular region of study for this project was the continental shelf off New Jersey where coastal upwelling fronts, riverine plumes, and internal solitary waves are common features. Our work is part of a large coordinated effort involving several other institutions (Dickey et al., 1999). The UCSB/OPL HyCODE mooring was deployed by OPL engineers, Derek Manov and Frank Spada, on May 16, 2000 on the New Jersey shelf (~24 m depth). This mooring was serviced and redeployed on July 25 and recovered on September 15, 2000. High temporal resolution measurements of IOPs and physical properties were collected at multiple depths. Hyperspectral instruments to measure AOPs were deployed in addition to the physical and IOP-optical instruments during the summer 2001 field program between June 20 and August 7, 2001. The data were processed and analyzed by OPL researchers Songnian Jiang and Grace Chang. Collaborative efforts have involved other HyCODE investigators, e.g., Oscar Schofield and Scott Glenn (Rutgers University), Mark Moline (Cal Poly), Emmanuel Boss (University of Maine), Scott Pegau (Oregon State University), and Alan Weidemann (NRL, Stennis). Additional details may be found in Dickey et al. (1999) and papers by Chang et al. (2002) and Chang and Dickey (2002) and on the web site: <http://www.opl.ucsb.edu/hycodeopl.html>.

WORK COMPLETED

We completed deployment and recovery of the first, second, and third HyCODE mooring deployments spanning the periods of May 16 - July 25, 2000; July 25 - September 15, 2000; and June 20 – August 7, 2001, respectively. All data have been processed. Datasets were complete with little to no bio-fouling seen in optical signals (see data reports by Chang et al., 2000; Dickey et al., 2000). Summer 2000 data and data reports have been made available to other HyCODE investigators via a CD-ROM.

A paper describing some of the results from the 2000 field observations is in press in the *Journal of Geophysical Research* (Chang et al., 2002). Mid-shelf mooring and tripod data and nearshore node data were analyzed statistically (frequency autospectra, coherence, autocorrelations, etc.) to investigate the temporal and spatial variability of hydrographic, physical, biological, and optical properties on scales of minutes to months and meters to ~50 km and to examine differences between nearshore and mid-shelf processes. Another paper focusing on solar transmission and radiant heating rates (Chang and Dickey, 2002) is about to be submitted as well. This paper utilizes HyCODE optical and physical data sets to explore the influences of cloud cover, chlorophyll and colored dissolved organic matter (CDOM) upon the magnitude and variability of solar transmission, sea surface albedo and radiant heating rates in the shallow coastal waters of the LEO-15 shelf region. These results are quite unique because of the concurrent, high temporal resolution optical and physical measurements in shallow waters and utilization of radiative transfer simulations.

RESULTS

The major processes contributing to bio-optical variability in summer 2000 were identified: e.g., coastal jet, upwelling front, tides, river flows, and internal solitary waves (Chang et al., 2002). Temporal decorrelation scales of optical and biological properties increased from nearshore (~1 day) toward the mid-shelf (2-3 days), whereas decorrelation scales for hydrographic properties were 2-3 days at both locations (Chang et al., 2002). Absorption at the mid-shelf location was dominated by phytoplankton and colored dissolved organic matter (CDOM), each accounting for roughly 50% of all absorbing materials at 440 nm. On the other hand, nearshore absorption was mainly influenced by particulate material (~70% of absorbing material) as compared to CDOM (~30% of absorbing material). Phytoplankton dominated the turbidity near the surface and intermediate depths and detritus dominated near the bottom. The interaction of tidal currents with the mean currents and the water mass/turbidity front were important for the formation of small-scale convergence and divergence zones (on the order of a few km) in the HyCODE experimental region. Frequency autospectra revealed that the M2 semidiurnal tides dominated temporal variability of physical, hydrographic, optical, and biological properties in both the nearshore and mid-shelf regions (Chang et al., 2002).

Time series of remote sensing reflectance were generated utilizing the optical model Hydrolight 4.0 (Mobley, 1994) and UCSB OPL mooring IOP data. Analyses of Hydrolight-generated $R_{rs}(\lambda)$ time series show that low salinity water masses in spring 2000, likely from the Hudson River, greatly influenced the optical properties at the HyCODE site. Correlations between $a_{t-w}(412):a_{t-w}(676)$, $a_{t-w}(676)-a_{t-w}(650)$ (a proxy for phytoplankton), and $b_{t-w}(412)$ versus the reciprocal of salinity (salinity^{-1}) show that these river flows resulted in increased CDOM in near-surface water. Spectral shifts of $R_{rs}(\lambda)$ were detected as well; the ratio of $R_{rs}(405):R_{rs}(675)$ increased from ~1.25 during times of particulate-dominated waters to >2 during times of CDOM-dominated near-surface waters.

Closure analyses between Hydrolight-generated (Mobley, 1994), and TSRB- and profiled radiometer-measured and calculated (using the relationship: $k_u = (-1/\Delta z) * \ln(L_u(\lambda)_2/L_u(\lambda)_1)$) upwelling and water-leaving radiance ($L_u(\lambda)$ and $L_w(\lambda)$, respectively) were performed with summer 2000 HyCODE data (courtesy of E. Boss and W. S. Pegau). These closure analyses resulted in r^2 of >0.95 and percent errors of less than 25% on average between measurement methods for $L_w(\lambda)$. Results from these analyses will prove useful for groundtruthing of remotely sensed data.

The results of the solar transmission/radiant heating rate study (Chang and Dickey, 2002) suggest that CDOM and possibly detritus can have significant impacts on the transmission of solar radiation through the water column in coastal waters and that the heat content of the upper layer can increase by orders of magnitude during periods of deep mixing.

IMPACT/APPLICATION

It is anticipated that results of the project will lead to 1) improved understanding of variability of inherent and apparent optical properties (IOPs and AOPs) and their relationships to each other as well as to physical processes on the ocean's continental shelves, 2) expanded usefulness and utility of high spectral and spatial resolution remotely sensed ocean color data, and 3) more accurate predictive interdisciplinary models of the coastal ocean; particularly primary productivity and optically-mediated radiant heating rate (inclusion of penetrative component of solar radiation) models in shallow waters.

All three of these points are fundamental to tactical naval applications in the coastal ocean and central to the ONR HyCODE program.

TRANSITIONS

Results from our statistical time series analyses will facilitate the development of interdisciplinary models (e.g., to predict the movement and distribution of biological matter, and study the anthropogenic effects on the coastal ocean) and facilitate the development and testing of ocean color algorithms to derive organic matter and primary production from remotely sensed data. Our data sets will also be valuable for development of a variety of optical, biological, and physical models and their couplings. Results of our work (see impacts above) should also be of interest to several levels of the operational Navy, particularly naval operations in the littoral zone.

RELATED PROJECTS

This project builds on the ONR Coastal Mixing and Optics (CMO) and PRIMER programs (see. <http://www.opl.ucsb.edu/cmo.html>). Results of our CMO work have already appeared in the reviewed literature (JGR, volume 106, number C5) as well as reports and conference proceedings (please see previous CMO Annual Report, 2001). Our HyCODE activity involves close collaborations with several other HyCODE, Naval Research Laboratory (NRL), and LEO-15 scientists. Our activity was also coordinated with the NRL COJET program. Hydrographic data from COJET are being used in analyses for interpretation.

REFERENCES

Chang, G. and T. Dickey, Solar transmission and radiant heating in coastal waters: optical influences, to be submitted to Journal of Geophysical Research.

Chang, G. C., T. D. Dickey, O. M. Schofield, A. D. Weidemann, E. Boss, W. S. Pegau, M. A. Moline, and S. M. Glenn, Nearshore physical processes and bio-optical properties in the New York Bight, J. Geophys. Res., in press, 2002.

Chang, G., S. Jiang, X. Yu, S. Zedler, D. Manov, D. Sigurdson, F. Spada, and T. Dickey, High resolution time series measurements of bio-optical and physical variability in the coastal ocean as part of HyCODE, Data Report: Hyperspectral Coupled Ocean Dynamics Experiment (HyCODE) Deployment II: 25 July – 15 September, 2000, Ocean Physics Laboratory Technical Report OPL-05-00, 2000.

Dickey, T., S. Ackleson, R. Arnone, P. Bissett, J. Donovan, S. Glenn, W. Hou, W. McBride, O. Schofield, J. Smart, and W. Snyder, Report of the First HyCODE Data Management and Distribution Workshop, Rutgers University, April 12-13, 1999.

Dickey, T., X. Yu, S. Jiang, S. Zedler, D. Manov, D. Sigurdson, F. Spada, and G. Chang, High resolution time series measurements of bio-optical and physical variability in the coastal ocean as part of HyCODE, Data Report: Hyperspectral Coupled Ocean Dynamics Experiment (HyCODE) Deployment I: 16 May – 25 July, 2000, Ocean Physics Laboratory Technical Report OPL-04-00, 2000.

Mobley, C. D. (1994) *Light and Water*, Academic Press, San Diego, CA, 592pp.

Web sites: <http://www.opl.ucsb.edu/hycodeopl.html> ; <http://www.opl.ucsb.edu/hycode.html>

PUBLICATIONS (2001-present)

Chang, G.C., T.D. Dickey, and A.J. Williams III, 2001, Sediment resuspension over a continental shelf during hurricanes Edouard and Hortense, *J. Geophys. Res.*, 106, 9517-9532.

Dickey, T.D. and A.J. Williams III, 2001, Interdisciplinary ocean process studies on the New England shelf, *J. Geophys. Res.*, 106, 9427-9434.

Kinkade, C.S., J. Marra, T.D. Dickey, and R. Weller, An annual cycle of phytoplankton biomass in the Arabian Sea, 1994-1995, 2001, as determined by moored optical sensors, *Deep-Sea Res. II*, 48, 1285-1301.

Dickey, T., S. Zedler, X. Yu, S.C. Doney, D. Frye, H. Jannasch, D. Manov, D. Sigurdson, J.D. McNeil, L. Dobeck, T. Gilboy, C. Bravo, D.A. Siegel, and N. Nelson, 2001, Physical and biogeochemical variability from hours to years at the Bermuda Testbed Mooring site: June 1994 – March 1998, *Deep-Sea Res. II*, 48 (8-9), 2105-2140.

Chang, G., T. Dickey, P. Bissett, and O. Schofield, 2001, High temporal resolution optical and physical time series data: Coastal Mixing and Optics and LEO-15, *Ocean Optics*, XV.

Dickey, T.D. and G.C. Chang, 2001, Recent advances and future visions: temporal variability of optical and bio-optical properties of the ocean, *Oceanography*, 14(3), 15-29.

Dickey, T.D., 2001. New technologies and their roles in advancing biogeochemical research, *Oceanography*, 14(4), 108-120.

Dickey, T., 2001, Sensors: inherent and apparent optical properties, *Encyclopedia of Ocean Science*, eds. J.H. Steele, S.A. Thorpe, and K.K. Turekian, Academic Press, London, 1313-1323.

Dickey, T., 2001, Emerging interdisciplinary sensors and systems for observing the world ocean, *Sea Tech.*, 42(12), 10-16.

Griffiths, G., R. Davis, C. Eriksen, D. Frye, P. Marchand, T. Dickey, and R. Weller, 2001, Towards new platform technology for sustained observations, in *Observing the Oceans in the 21st Century*, eds. C.J. Koblinksky and N.R. Smith, Australian Bureau of Meteorology, 324-338.

Send, U., R. Weller, S. Cunningham, C. Eriksen, T. Dickey, M. Kawabe, R. Lukas, M. McCartney, and S. Osterhuis, 2001, Oceanographic time series observatories, *Observing the Ocean for Climate in the 21st Century*, eds. C.J. Koblinksky and N.R. Smith, Australian Bureau of Meteorology, 376-390.

Dickey, T. and P. Falkowski, 2002, Solar energy and its biological-physical interaction in the sea, eds. A.R. Robinson, J.J. McCarthy, and B.J. Rothschild, Chapter 10, *The Sea*, Vol. 12, John Wiley, New York, 401-440.

Souza, A.J., T.D. Dickey, and G.C. Chang, 2002, Modeling water column structure and suspended particulate matter in the Middle Atlantic continental shelf during passage of Hurricanes Edouard and Hortense, *J. Mar. Res.*, 59, 819-842.

Jones, B., M. Noble, and T. Dickey, 2002, Hydrographic and particle distributions over the Palos Verdes continental shelf: spatial, seasonal, and tidal variability, *Cont. Shelf Res.*, 22(6-7), 945-965.

Dickey, T., 2002, A vision of oceanographic instrumentation and technologies in the early 21st century, Chapter 9, *Oceans 2020: Science for Future Needs*, eds. J.G. Field, G. Hemphill, and C.P. Summerhayes, Island Press, Washington, DC, 213-256.

Weller, R., A.S. Fischer, D.L. Rudnick, C.E. Eriksen, T. Dickey, J. Marra, C. Fox, and R. Leben, 2002. Moored observations of upper ocean response to the monsoons in the Arabian Sea during 1994-1995, *Deep-Sea Res. II*, 49, 2195-2230.

Chang, G.C., T.D. Dickey, O. Schofield, A.D. Weidemann, E. Boss, W.S. Pegau, M. Moline, and S.M. Glenn, 2002. Temporal and spatial scales of physical and bio-optical variability on the New York Bight inner continental shelf, *J. Geophys. Res.*, in press.

Zedler, S.E., T.D. Dickey, S.C. Doney, J.F. Price, X. Yu, and G.L. Mellor, 2002, Analysis and simulations of the upper ocean's response to Hurricane Felix at the Bermuda Testbed Mooring site: August 13 – 23, 1995, *J. Geophys. Res.*, in press.

Yu, X., T. Dickey, J. Bellingham, D. Manov, and K. Streilein, 2002, The application of autonomous underwater vehicles for interdisciplinary measurements in Massachusetts and Cape Cod Bays, *Cont. Shelf Res.*, in press.

Zheng, X., T. Dickey, and G. Chang, 2002, Variability of the downwelling diffuse attenuation coefficient with consideration of inelastic scattering, *Appl. Optics*, in press.

Dickey, T., 2002, Emerging ocean observations for interdisciplinary data assimilation systems, *J. Mar. Sys.*, in press.